

# MITEARREST<sup>®</sup>

INSECTICIDAL BEDDING FOR LABORATORY RODENTS

## For Laboratory, Commercial or Veterinary Use

EPA Reg. No. 56783-2

EPA Est. No. 5693 MA 1

Permethrin impregnated cotton bedding to control mite infestations on laboratory rodents.

### ACTIVE INGREDIENT:

+\* Permethrin ----- 7.4%

INERT INGREDIENTS: ----- 92.6%

-----  
Total 100.0%

+ (3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2 dichloroethenyl)2, 2-dimethylcyclopropanecarboxylate.

\* Cis/trans ratio: Min. 35% (+/-) cis and max. 65% (+/-) trans.

## CAUTION: KEEP OUT OF REACH OF CHILDREN

### FIRST AID

#### INGESTION - If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

#### EXPOSURE, SKIN IRRITATION - If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### EYES - If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

# MITEARREST<sup>®</sup>

INSECTICIDAL BEDDING FOR LABORATORY RODENTS

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**APPLICATION INSTRUCTIONS:** Place MITEARREST bedding directly in rodent cage to be treated. Use 2-3 MITEARREST balls per week for each rodent. Avoid wetting the MITEARREST bedding; to prevent cage flooding, care should be taken to place MITEARREST bedding away from the water bottle spout. Replace soiled MITEARREST bedding; it is recommended that fresh MITEARREST bedding be added weekly, at the time when other bedding is changed. Continue using the product for four (4) to six (6) weeks, or until mites or their eggs are no longer present in the fur of rodent. As mite eggs are not affected by this treatment, extended use of the product (until all eggs have hatched or are no longer present) will provide greater long-term control and prevent reinfestation.

**STORAGE:** Store at room temperature. Do not contaminate water, food or feed by storage.

### DISPOSAL

#### If empty:

Do not reuse this container.  
Place in trash or offer for recycling if available.

#### If partly filled:

Call your local solid waste agency or  
1-800-234-8425 for disposal instructions.  
Never place unused product down any indoor or  
outdoor drain.

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS & DOMESTIC ANIMALS:** Avoid contact with eyes and skin. Contact with skin may result in temporary numbing and tingling sensations. Wash thoroughly with soap after handling.

**ENVIRONMENTAL HAZARDS:** This product is extremely toxic to fish and other aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of wastes or equipment washwaters.

**NOTE:** This product contains a pyrethroid insecticide which the caged animal may ingest or come into dermal contact with. Researchers should consider the possible effects of contact with this insecticide before using the product.

U.S. Patent No. 4,662,104

© 2004, ECOHEALTH, INC.

33 MT. VERNON ST. BOSTON, MASSACHUSETTS 02108

617-742-2400 WWW.MITEARREST.COM

Net Weight 2.5 lbs.